

THE RELATIONSHIP BETWEEN CHLOROPHYLL- α AND TOTAL PHOSPHORUS (TOTAL-P) IN PERUPUK LAKE, PERHENTIAN RAJA SUB-DISTRICT, KAMPAR, RIAU

Andika Ramadhani^{1*}, Asmika Harnalin Simarmata¹, Budijono¹

¹Department of Aquatic Resources Management, Faculty of Fisheries and Marine, Universitas Riau, Pekanbaru, 28293 Indonesia

*andika.ramadhani4548@student.unri.ac.id

ABSTRACT

The province of Riau has many freshwater bodies that play a very important role in the lives of those around them. This study aimed to determine the relationship between chlorophyll a and Total-P (Total Phosphorus) concentrations and to assess the trophic status of Perupuk Lake in Perhentian Raja Sub-District, Kampar Regency, Riau Province. The research was conducted from October to December 2024 at three stations: Station 1 (inlet, no settlement), Station 2 (lake main basin with fish cages), and Station 3 (outlet with settlement and agricultural activity). Sampling was performed 5 times at 1-week intervals at the surface and at 2 Secchi depths. Parameters measured included chlorophyll a, Total-P, temperature, transparency (Secchi depth), pH, dissolved oxygen (DO), free carbon dioxide (CO₂), and nitrate. Results showed that chlorophyll a concentration ranged from 11.04 to 14.14 $\mu\text{g/L}$, with the highest at Station 2 (14.14 $\mu\text{g/L}$) and the lowest at Station 1 (11.04 $\mu\text{g/L}$). Total-P concentrations ranged from 0.0168–0.0338 mg/L, following the same pattern: highest at Station 3 (0.0338 mg/L) and lowest at Station 1 (0.0168 mg/L). Simple regression analysis showed a non-significant relationship between Total-P and chlorophyll-a ($R^2 = 0.0266\text{--}0.0774$), indicating that, in mesotrophic waters, other factors, such as light and nitrogen availability, also influence chlorophyll-a concentration. Based on chlorophyll a and Total-P concentrations, Perupuk Lake is categorized as a mesotrophic water body, with supporting parameters within optimal ranges for aquatic organisms.

Keywords: Chlorophyll-a, Total-P, Mesotrophic, Oxbow Lake, Water Quality

1. INTRODUCTION

The province of Riau has many freshwater bodies that play a very important role in the lives of those around them. Kampar Regency is one of the regencies in Riau Province with an area of 10,928.20 km². Public water bodies in Kampar Regency include rivers, reservoirs, and oxbow lakes. An oxbow lake is a lake that forms when a river changes course due to erosion and sedimentation along its flow¹. Lake Perupuk is an oxbow lake located in Kampung Pinang Village, Perhentian Raja Subdistrict, Kampar Regency, Riau Province. This lake covers approximately

4.15 ha, has an average depth of 5 m, and was formed when the flow of the Kampar River was interrupted. Lake Perupuk has important ecological, socio-cultural, and economic functions. Ecologically, this lake is a habitat for aquatic organisms and maintains hydrological balance. From a socio-economic perspective, the lake is used by the surrounding community for fishing and for tourism development².

Various activities are carried out by the community around Lake Perupuk, including residential settlements, cattle farming, oil palm plantations, and fish farming. These activities have the potential

to contribute organic and inorganic materials to the water. Organic inputs undergo decomposition into nutrients such as nitrogen and phosphorus. Increased concentrations of these nutrients can stimulate phytoplankton growth, thereby increasing chlorophyll a concentration in the water³. Chlorophyll a is a green pigment in plant cells that plays an important role in photosynthesis in water. Chlorophyll a can be used as an indicator of phytoplankton biomass and water fertility⁴. Stated that an increase in chlorophyll a concentration in a body of water is related to the phosphate concentration in that water. Phosphorus is a macronutrient that is very important for the survival of aquatic organisms because it plays a role in the storage and transfer of energy within cells.

Research on the relationship between chlorophyll a and Total-P has been conducted in several places. Hutajulu⁵ reported that chlorophyll a and total phosphorus have a very strong relationship in the Koto Panjang Hydroelectric Power Plant Reservoir. However, similar research has never been conducted in Lake Perupuk, which differs from other oxbow lakes due to agricultural and residential activities around it. Therefore, this study is important for providing a comprehensive overview of the relationship between water fertility and chlorophyll a in Lake Perupuk and for serving as a basis for sustainable water resource management.

The objectives of this study are to determine the relationship between chlorophyll a and total phosphorus in Lake Perupuk and to assess the trophic status of the water. The benefit of this study is that, by knowing the concentration of Total-P, the concentration of chlorophyll a in Lake Perupuk can be estimated, which can then serve as a basis for future water management

2. RESEARCH METHOD

This and Place

The research was conducted from October to December 2024 in the waters of Lake Perupuk, Kampung Pinang Village,

Perhentian Raja Subdistrict, Kampar Regency, Riau Province. Field observations and measurements of several parameters were conducted at the research site. At the same time, laboratory analysis was performed at the Water Productivity Laboratory of the Faculty of Fisheries and Marine Sciences, Universitas Riau.




Figure 1. Research location

Method

The methods used in this study included surveys, direct observation at the research site, and water sampling and water quality measurements. The data collected included primary and secondary data. Primary data were obtained from direct water-quality measurements or from laboratory analyses. Meanwhile, secondary data consisted of topographical data for Kampung Pinang Village, Perhentian Raja Subdistrict, Kampar Regency, obtained from the literature.

Procedures

Determination of the Research Station

Research stations were determined using purposive sampling based on location characteristics. Three research stations were established as follows: Station I: This is the water inlet to the lake. There are no residential areas around this station, but there are oil palm plantations (0°21'55"N 101°27'02"E). Station II: This is the central part of the lake with fish farming and oil palm plantations (0°21'53"N 101°27'10"E). Station III: This is the lake's water outlet with residential settlements, fish farming, and oil palm plantations (0°22'12"N 101°27'13"E).

Water Sampling Chlorophyll-a

The study used a cross-sectional survey with a sampling design. Sampling was conducted five times at one-week intervals at two depths: surface and 2 Secchi depths. Water samples were collected using a Kemmerer water sampler for the 2 Secchi depths and directly from the surface for surface samples. A total of 500 mL of water samples were taken and covered with aluminum foil. In the laboratory, the samples were filtered using Millipore paper under vacuum. Millipore paper containing chlorophyll a was ground with 90% acetone solution (5 mL + 3.5 mL), then centrifuged at 2000 rpm for 10 minutes. The absorbance of the supernatant was measured at 665 nm and 750 nm using a spectrophotometer. The chlorophyll-a concentration was calculated using the formula by Agung et al.⁴:

$$\text{Klorofil-a } (\mu\text{g/L}) = 11,9 (A_{665} - A_{750}) \frac{V}{L} \times \frac{1000}{S}$$

Description:

A665 = Spectrophotometer absorption
 λ 665

A750 = Spectrophotometer absorption
 λ 750

V = Volume of acetone extract
used (mL)

S = Volume of filtered sample
(mL)

L = Light path length or cuvette
width (1cm)

11.9 = Constant (constancy)

1000 = Conversion from L to mL

Total-P

A 25-mL sample was treated with 30% H₂SO₄ and 5% K₂S₂O₈, heated to 100°C, and cooled to 60°C. A pp indicator was added, and the solution was titrated with 10 N NaOH until a pink color appeared. The sample was diluted to 25 mL with distilled water; 5 mL was taken, and 0.2 mL of ammonium molybdate and 1 drop of SnCl₂ were added. Absorbance was measured at λ = 690 nm using the APHA standard curve. Supporting parameters included temperature measured with a thermometer, pH with a pH indicator, turbidity with a Secchi disk, DO using the Winkler method, free CO₂ by

titration with Na₂CO₃, and nitrate by spectrophotometer at λ 543 nm

3. RESULT AND DISCUSSION

Chlorophyll-a

The concentration of chlorophyll-a at the surface ranged from 11.04 to 14.08 $\mu\text{g/L}$, with the lowest concentration at Station 1 (11.04 $\mu\text{g/L}$) and the highest at Station 2 (14.08 $\mu\text{g/L}$). At a depth of 2 Secchi, the concentration ranged from 11.37 to 14.14 $\mu\text{g/L}$ (Figure 1). The vertical profile shows that chlorophyll a concentration in the water column are higher than at the surface, especially at Station 2. This is thought to be because nutrient concentrations in the water column are higher, resulting in more optimal photosynthesis³. The low chlorophyll a at the surface of Station 1 was due to low phosphate, CO₂, and brightness levels, which inhibited photosynthesis. The high chlorophyll a at Station 2 was associated with high CO₂ and nutrient concentrations. According to the classification of Agung et al.⁴ based on chlorophyll a, Lake Perupuk is mesotrophic (8–25 $\mu\text{g/L}$), as its average chlorophyll a is 11.40–14.14 $\mu\text{g/L}$.




Figure 2. Chlorophyll-a concentration in Lake Perupuk

Total-P

Concentrations at the surface ranged from 0.0168 to 0.0284 mg/L and in the water, column ranged from 0.0252 to 0.0338 mg/L (Figure 2). Total-P was highest at Station 3 (outlet with residential area) and lowest at Station 1 (inlet). Total-P concentrations in the water column were higher than at the surface at all stations. This is because phosphorus has a higher density than water, so it tends to settle in deeper

layers⁶. The high Total-P at Station 3 is associated with fish farming and oil palm cultivation in the surrounding area. During the rainy season, fertilizer residues and agricultural waste are carried into the water. The low Total-P at Station 1 is due to its position as an inlet without significant organic input. Based on the classification of Agung et al.⁴, with a Total-P of 0.01–0.035 mg/L, Lake Perupuk is classified as mesotrophic.




Figure 3. Total-P concentration in Lake Perupuk

Table 1. The relationship between chlorophyll-a and total-P

Station	Chlorophyll-a	Total-P	Regression Equation	R ²
1	X	Y	$y = 10,564 + 31,152x$	0,0774
2	X	Y	$Y = 13,036 + 37,664x$	0,0275
3	X	Y	$y = 12,656 + 0,4043\ln(x)$	0,0266

Based on a simple regression analysis, the relationship between Total-P concentration and chlorophyll-a obtained during research in Lake Perupuk was found to be insignificant⁴. This is because Lake Perupuk is in a mesotrophic state, with moderate phosphorus availability in the water, which is not a major limiting factor for chlorophyll-a formation. In eutrophic waters, phosphorus is no longer the main factor influencing the increase in chlorophyll-a. Therefore, the concentration of chlorophyll-a in Lake Perupuk is strongly suspected to be more influenced by other factors, such as other physical and chemical parameters.

Water Quality

Temperature measurements during the study at Lake Perupuk ranged from 28 to 30.2°C (Figure 4). Agung et al.⁴ stated that the optimal temperature range for the life and development of aquatic organisms is 25–32°C. The surface water temperature range recorded at Lake Perupuk during the study, namely 28–30.2°C, supports this opinion, indicating that the temperature during the study was still capable of supporting the life of aquatic organisms.




Figure 4. Vertical temperature profile in Lake Perupuk

Acidity Level (pH)

The results of acidity level (pH) measurements found during research in Lake Perupuk were 6 (Figure 5). Based on the pH values obtained, the waters of Lake Perupuk are acidic but can still support aquatic organisms. The acidity level during the study was the same at the surface and at a depth of 2 Secchi.

According to Effendi³, aquatic organisms generally live in waters with a pH of 5–9, with an optimal pH range of 6–8. The pH of Lake Perupuk, around 6, indicates that water conditions are still within acceptable limits and support aquatic life.

Dissolved Oxygen

The average dissolved oxygen concentration measured during the study ranged from 4.776 to 5.358 mg/L. The

dissolved oxygen concentration at the surface was relatively higher than in the water column. This is because light penetrates more deeply at the surface, enabling maximum photosynthesis. This is supported by Wetzel⁷, who states that photosynthesis by phytoplankton and aquatic plants produces dissolved oxygen in water. Based on the dissolved oxygen concentrations obtained during the study, it can still support the life of these aquatic organisms. This is in accordance with Ismail⁸, who states that the minimum content sufficient to support the normal life of aquatic organisms is 4 mg/L.




Figure 6. Vertical profile of dissolved oxygen in Lake Perupuk

Free Carbon Dioxide (CO₂)

The average free carbon dioxide concentration during the study ranged from 10.3 to 10.6 mg/L. The average free CO₂ results during the study still support the life of aquatic organisms in Lake Perupuk. This is in line with the opinion of Agung et al. in Lumbangaol⁹, who stated that the maximum free carbon dioxide content in water is 10-15 mg/L. According to Octasari et al.¹⁰, the ideal carbon dioxide (CO₂) level for the survival of aquatic organisms is approximately 15 mg/L. Levels above this are very dangerous because they inhibit oxygen (O₂) binding.

The average free carbon dioxide (CO₂) concentration measured during the study at Lake Perupuk ranged from 8 to 11 mg/L at all three observation stations. Vertically, CO₂ concentrations tended to increase with depth, especially in the lower water layers. This pattern was relatively similar across all stations, although there were differences in

concentration values at certain depths. The lower CO₂ concentration in the surface layer is thought to be related to the photosynthetic activity of phytoplankton and aquatic plants that utilize CO₂, as well as gas diffusion into the atmosphere. Conversely, the increase in CO₂ concentration in the deeper layers is caused by the respiration of aquatic organisms and more intensive decomposition of organic matter, as well as limited water mass mixing at that depth.




Figure 7. Vertical profile of free dissolved carbon dioxide in Lake Perupuk

Phosphate (PO₄³⁻)

The average orthophosphate concentration measurements during the study ranged from 0.0168 to 0.0338 mg/L at the three observation stations. The average phosphate concentration at Station 3 was higher than at the other stations. This is because Station 3 receives organic matter from oil palm plantations and fish farming activities in the surrounding area, which is subsequently decomposed into nutrients.




Figure 8. Vertical phosphate profile in Lake Perupuk

Effendi³ classified low water fertility levels as 0-0.02 mg/L, moderate water fertility levels as 0.02-0.05 mg/L, and high fertility levels as 0.051-0.1 mg/L, with very

fertile hypertrophic levels ranging from 0.101-0.2 mg/L. Based on these criteria, Lake Perupuk is classified as mesotrophic to eutrophic, with phosphate concentrations ranging from 0.0168 to 0.0338 mg/L

4. CONCLUSION

Based on research in Lake Perupuk, chlorophyll-a concentration ranged from 11.396 to 14.136 µg/L, with the highest value at Station II and the lowest at Station I. The high chlorophyll-a concentration at Station II was influenced by the abundance of phytoplankton driven by high total phosphorus (Total-P) and nitrate levels. The vertical profile showed that chlorophyll a concentration tended to be higher at a depth of 2 Secchi at stations with high brightness levels, allowing for more optimal light

penetration for photosynthesis. The relationship between chlorophyll a and Total-P follows a non-linear pattern, indicating that light availability is also an important factor in addition to nutrients in the chlorophyll formation process. In general, Lake Perupuk is categorized as mesotrophic based on chlorophyll-a levels. Other water quality parameters, such as temperature (28°C-30.2°C), pH (with an average of 6), dissolved oxygen (4.776-5.358 mg/L), and free carbon dioxide (10.45-10.608 mg/L), remain within ranges that support aquatic life. Therefore, the water conditions of Lake Perupuk remain productive and suitable for supporting phytoplankton growth, although regular monitoring is necessary to anticipate the possibility of increased eutrophication.

REFERENCES

1. Dian, D. *Distribusi Vertikal Klorofil-a di Danau Baru Desa Buluh Cina Kecamatan Siak Hulu Kabupaten Kampar, Riau*. Fakultas Perikanan dan Kelautan. Universitas Riau. Pekanbaru, 2013.
2. Siburian, R. *Pembangunan Pariwisata Berkelanjutan: Tantangan dan Peluang di Kawasan Danau Toba*. LIPI Press. Jakarta, 2020.
3. Effendi, H. *Telaah Kualitas Air Bagi Pengelolaan Sumber Daya dan Lingkungan Perairan*. Kanisius, Yogyakarta. 2003; 249p.
4. Agung, A., Zainuri, M., Wirasatriya, A., Maslukah, L., Subardjo, P., Suryosaputro, A.A.D., Handoyo, G.. Analisis Sebaran Klorofil-a dan Suhu Permukaan Laut sebagai Fishing Ground Potensial (Ikan Pelagis Kecil) di Perairan Kendal, Jawa Tengah. *Buletin Oseanografi Marina*, 2018; 7(2): 67-74
5. Hutajulu, O. *Hubungan Fosfat dengan Klorofil-a di Zona Lakustrin dan Transisi Waduk PLTA Koto Panjang Kabupaten Kampar Provinsi Riau*. Fakultas Perikanan dan Kelautan. Universitas Riau. Pekanbaru, 2014.
6. Walukow, A.F. Penentuan Status Mutu Air dengan Metode STORET di Danau Sentani, Jayapura, Provinsi Papua. *Berita Biologi*, 2010; 10(3): 277-281
7. Wetzel, R.G. *Limnology: Lake and River Ecosystems*. Gulf Professional Publishing. Academic Press. San Diego, 2001; 1006p.
8. Ismail, I. *Kajian Kualitas Air bagi Organisme Perairan*. Bogor: Fakultas Perikanan dan Ilmu Kelautan, Institut Pertanian Bogor, 1994.
9. Lumbangaol, S.M. *Studi Kualitas Air di Perairan Danau Baru Desa Buluh Cina Kecamatan Siak Hulu Kabupaten Kampar Provinsi Riau*. Fakultas Perikanan dan Kelautan. Universitas Riau. Pekanbaru, 2014.
10. Octasari, A., Agustiani, R., Susilo, H. Analisis Kualitas Air pada Budidaya Ikan di Karamba Jaring Apung Sungai Musi Kota Palembang. *Jurnal Ilmu-Ilmu Perikanan dan Budidaya Perairan*, 2018; 13(1): 31-38